

# PEDIATRIC NAUSEA/VOMITING

HPI	Signs & Symptoms	Differential Diagnosis
<ul style="list-style-type: none"> <li>• Time of last meal</li> <li>• Last bowel movement</li> <li>• Last emesis</li> <li>• Relation to food</li> <li>• Menstrual cycle</li> <li>• Trauma</li> </ul>	<ul style="list-style-type: none"> <li>• Pain, character</li> <li>• Distention</li> <li>• Constipation</li> <li>• Diarrhea</li> <li>• Radiation</li> </ul>	<ul style="list-style-type: none"> <li>• Cardiac</li> <li>• Drugs (NSAID's, antibiotics, narcotics)</li> <li>• GI or renal disorders</li> <li>• Food or toxin induced</li> <li>• Pregnancy, GYN</li> </ul>

	EMR	EMT	PM
1. Provide <a href="#">Pediatric Routine Medical Care</a> - treat life threatening problems with airway, breathing circulation and disability.	•	•	•
2. Obtain ECG, monitor Capnography.		•	•
3. <b><a href="#">ONDANSETRON</a>:</b> ≥ 10kg and <40 0.1 mg/kg IVP/IO, over 30 sec., max. 4 mg ≥40 kg: 4 mg IVP/IO, over 30 sec.			•
or 4 mg ORAL, may repeat 10 min., max. 8 m	•		•

**NOTE:**

1. Do not administer to patients under 10kg
2. Do not administer to patients who are pregnant

**Use Calculator for dosing**